arXiv:1609.05627v1 [math.AG] 19 Sep 2016

ALGEBRAIC CYCLES ON THE FANO VARIETY OF LINES OF A CUBIC FOURFOLD

KALYAN BANERJEE

ABSTRACT. In this text we try to find an obstruction to rationality of a cubic fourfold by studying the zero cycles on the Fano variety of lines of the cubic.

1. INTRODUCTION

In the article [BG] the authors were discussing the kernel of the push-forward homomorphism at the level of algebraically trivial one cycles modulo rational equivalence, induced by the closed embedding of a smooth hyperplane section of a cubic fourfold into the cubic itself. It has been proved that the kernel of this homomorphism is countable when we consider a very general hyperplane section. It was a natural question at that time that can we formulate the rationality problem of a very general cubic fourfold in terms of this kernel. That is what is the obstruction to rationality in terms of the kernel.

In [GS][theorem 7.5] it has been proved that if the ground field k is such that the class of the affine line is not a zero divisor in the Grothendieck ring of varieties over k and if the cubic fourfold is rational then the Fano variety of lines on the cubic fourfold is birational to the Hilbert scheme of two points on a K3 surface. This is the starting points of this paper. We start with a birational map from the Hilbert scheme of two points on a K3 surface to theh Fano variety of lines and try to formulate the rationality question in terms of zero cycles on the Fano variety. It is also known, due to Hassett [Ha] that for a certain class of rational cubic fourfolds the Hilbert scheme is actually isomorphic to the Fano variety of lines.

To proceed to find the obstruction to rationality in terms of the zero cycles on the Fano variety of lines, we study algebraic cycles on symmetric powers of a smooth projective variety following Collino [Collino], where the algebraic cycles on symmetric powers of a smooth projective curve was studied. It has been proved in [Collino] that the closed embedding of

one symmetric power of a smooth projective curve into a bigger symmetric power of the same curve induces injective push-forward at the level of Chow groups. We follow this approach and try to understand the algebraic cycles on the blowup of $\mathrm{Sym}^2 S$ along the diagonal where S is a smooth projective surface. This blowup is the Hilbert scheme of two points on a smooth projective surface. Following Collino's approach we study a correspondence between the Hilbert scheme of two points on a surface and the strict transform the surface itself under the blowup along the diagonal. Our main result of this paper is the following:

Let X be a smooth cubic fourfold in $\mathbb{P}^5$ and let $F(X)$ be the Fano variety of lines on the cubic fourfold X. If the cubic X is rational and if the indeterminacy locus of the rational map $\mathrm{Hilb}^2 S \to F(X)$ is smooth, then there exists a projective curve, or a projective surface (admitting a finite map from a ruled surface) in $F(X)$ such that the kernel of the push-forward (induced by the closed embedding of the curve or the surface inside $F(X)$) at the level of Chow groups of zero cycles in torsion. Contra-positively we can say that if for all projective curves and projective surfaces in $F(X)$, the kernel of the push-forward induced by the closed embedding of the curve or the surface has non-torsion elements, then the cubic is non-rational.

2. Algebraic cycles on Hilbert schemes

In this section we are going to prove the following. Let S be a smooth, projective algebraic surface. Consider the diagonal embedding of S into $\mathrm{Sym}^2 S$. Then the blow up of $\mathrm{Sym}^2 S$ along S is the Hilbert scheme of two points on S. Let $\widetilde{\mathrm{Sym}^2 S}$ denote this Hilbert scheme and let $\widetilde{S}$ denote the inverse image of S in $\widetilde{\mathrm{Sym}^2 S}$ (This copy of S is different from the image of the diagonal embedding and it is the image of the closed embedding $s \mapsto [s, p]$). Then the embedding of $\widetilde{S}$ into $\widetilde{\mathrm{Sym}^2 S}$ induces an injective push-forward homomorphism at the level of Chow group of zero cycles.

To prove that we prove that if blow up $\mathrm{Sym}^n X$ along a subvariety Z, which does not contain a copy of $\mathrm{Sym}^m X$ for $m \leq n$, then the closed embedding of the inverse image of $\mathrm{Sym}^m X$ into the blow up of $\mathrm{Sym}^n X$ induces injective push-forward homomorphism at the level of Chow groups of zero cycles.

Theorem 2.1. *Let X be a smooth projective variety. Let $\widetilde{\mathrm{Sym}^n X}$ be the blow up of $\mathrm{Sym}^n X$ along a smooth subvariety Z, such that the blow-up will be a*

smooth projective variety. Let $\mathrm{Sym}^m X$ *intersect* Z *transversally. Then the closed embedding of the strict transform of* $\mathrm{Sym}^m X$ *into* $\widetilde{\mathrm{Sym}^n X}$ *induces an injective push-forward homomorphism at the level of Chow groups of zero cycles.*

Proof. To prove the theorem we follow the approach as in [Collino]. First consider the correspondence Γ on $\mathrm{Sym}^n X \times \mathrm{Sym}^m X$ given by

$$(\pi_n \times \pi_m)(Graph(pr))$$

where pr is the projection from X^n to X^m and π_i is the natural quotient morphism from X^i to $\mathrm{Sym}^i X$. Let f be the natural morphism from $\widetilde{\mathrm{Sym}^n X}$ to $\mathrm{Sym}^n X$ and f' be the restriction of f to $\widetilde{\mathrm{Sym}^m X}$, which is the inverse image of $\mathrm{Sym}^m X$. Consider the correspondence Γ' given by $(f \times f')^*(\Gamma)$. Then a computation following [Collino] shows that $\Gamma'_* j_*$ is induced by $(j \times id)^*(\Gamma')$, where j is the closed embedding of $\widetilde{\mathrm{Sym}^m X}$ into $\widetilde{\mathrm{Sym}^n X}$. Consider the following diagram.

$$\begin{array}{ccc} \widetilde{Sym^m X} \times \widetilde{\mathrm{Sym}^m X} & \xrightarrow{j \times id} & \widetilde{\mathrm{Sym}^n X} \times \widetilde{\mathrm{Sym}^m X} \\ \downarrow{\scriptstyle f' \times f'} & & \downarrow{\scriptstyle f \times f'} \\ \mathrm{Sym}^m X \times \mathrm{Sym}^m X & \xrightarrow{i \times id} & \mathrm{Sym}^n X \times \mathrm{Sym}^m X \end{array}$$

This diagram gives us the following at the level of algebraic cycles.

$$(j \times id)^*(f \times f')^*(\Gamma) = (f' \times f')^*(i \times id)^*(\Gamma)$$

that is equal to

$$(f' \times f')^*(\Delta + Y)$$

by [Collino], where Y is supported on $\mathrm{Sym}^m X \times \mathrm{Sym}^{m-1} X$, and Δ is the diagonal of $\mathrm{Sym}^m X \times \mathrm{Sym}^m X$. Now

$$(f' \times f')^*(\Delta) = \Delta + E_1$$

where

$$E_1 \subset (E \times E) \cap (\widetilde{\mathrm{Sym}^m X} \times \widetilde{\mathrm{Sym}^m X})$$

$E$is the exceptional locus, that is the pre-image of Z. So then by the Chow moving lemma, we can take the support of a zero cycle away from $E \cap \widetilde{\mathrm{Sym}^m X}$, which is a proper closed subscheme of $\widetilde{\mathrm{Sym}^m X}$, and we get

$$\rho^* \Gamma'_* j_*(z) = \rho^*(z + z_1) = \rho^*(z)$$

where ρ is the open embedding of the complement of $\widetilde{\mathrm{Sym}^{m-1}X}$ in $\widetilde{\mathrm{Sym}^{m}X}$, and z_1 is supported on $\widetilde{\mathrm{Sym}^{m-1}X}$. Now consider the following commutative diagram with the rows exact.

$$\begin{array}{ccccc} CH^*(\widetilde{\mathrm{Sym}^{m-1}X}) & \xrightarrow{j'_*} & CH^*(\widetilde{\mathrm{Sym}^{m}X}) & \xrightarrow{\rho^*} & CH^*(U) \\ \downarrow & & \downarrow{\scriptstyle j_*} & & \downarrow \\ CH^*(\widetilde{\mathrm{Sym}^{m-1}X}) & \xrightarrow{j''_*} & CH^*(\widetilde{\mathrm{Sym}^{n}X}) & \longrightarrow & CH^*(V) \end{array}$$

Suppose that $j_*(z) = 0$, by the above discussion we have $\rho^*\Gamma'_* j_*(z) = \rho^*(z) = 0$. Hence by the localisation exact sequence there exists z' such that $j'_*(z') = z$. So by induction $CH_*(\widetilde{\mathrm{Sym}^{m-1}X})$ to $CH_*(\widetilde{\mathrm{Sym}^{n}X})$ is injective.

So we have $z' = 0$ and hence $z = 0$. So we have the theorem.

□

Let S be a smooth projective algebraic surface. Let us consider the diagonal embedding of S into $\mathrm{Sym}^2 S$ and the embedding of S into $\mathrm{Sym}^2 S$ given by $s \mapsto [s, p]$, where p is a fixed closed point on S. Let $\widetilde{\mathrm{Sym}^2 S}$ be the blow up of $\mathrm{Sym}^2 S$ along the image of the diagonal embedding and let $\widetilde{S}$ be the strict transform of the other copy of S prescribed by the late embedding.

Corollary 2.2. *The closed embedding of $\widetilde{S}$ into $\widetilde{\mathrm{Sym}^2 S}$ induces injective push-forward homomorphism at the level of Chow groups of zero cycles.*

Proof. Follows from 2.1. □

$\widetilde{\mathrm{Sym}^2 S}$ is nothing but the Hilbert scheme of two points on S. The above corollary tell us in particular that the closed embedding of $\widetilde{S}$ into the Hilbert scheme of two points on a surface S induces injective push-forward homomorphism at the level of Chow group of zero cycles.

Using the same technique we can prove that if we blow up the Hilbert scheme once again and consider the total transform of the surface $\widetilde{S}$, then the closed embedding of the total transform in the blow-up gives rise to an injective push-forward homomorphism at the level of Chow group of zero cycles.

3. Blow down of symmetric powers

In this section we are going to consider blow downs of $\mathrm{Sym}^n X$, for a smooth projective variety X, and consider the images of $\mathrm{Sym}^m X$ under the blow down. Then we want to investigate the nature of the push-forward homomorphism induced at the level of Chow groups by the closed embedding of the blow down of $\mathrm{Sym}^m X$ into the blow down of $\mathrm{Sym}^n X$.

Theorem 3.1. *Let* $\widetilde{\mathrm{Sym}^n X}$ *be a smooth blow down of* $\mathrm{Sym}^n X$ *and let* $\widetilde{\mathrm{Sym}^m X}$ *denote the image of* $\mathrm{Sym}^m X$ *under this blow down. Then the closed embedding of* $\widetilde{\mathrm{Sym}^m X}$ *into* $\widetilde{\mathrm{Sym}^n X}$ *induces an injective push-forward homomorphism at the level of Chow groups.*

Proof. As before consider the correspondence Γ given by $(\pi_n \times \pi_m)(Graph(pr))$, where pr is the projection from X^n to X^m and π_i is the natural morphism from X^i to $\mathrm{Sym}^i X$. Let $\widetilde{f}$ denote the morphism from $\mathrm{Sym}^n X$ to $\widetilde{\mathrm{Sym}^n X}$, and f' its restriction to $\widetilde{\mathrm{Sym}^m X}$. Consider the correspondence Γ' on $\widetilde{\mathrm{Sym}^n X} \times \widetilde{\mathrm{Sym}^m X}$ given by $(f \times f')_*(\Gamma)$. Then following [Collino], we have that, $\Gamma'_* j_*$ is induced by $(j \times id)^*(\Gamma')$, where j is the closed embedding of $\widetilde{Sym^m X}$ into $\widetilde{\mathrm{Sym}^n X}$. Now consider the following Cartesian square.

$$\begin{array}{ccc} Sym^m X \times \mathrm{Sym}^m X & \xrightarrow{i \times id} & \mathrm{Sym}^n X \times \mathrm{Sym}^m X \\ \downarrow{\scriptstyle f' \times f'} & & \downarrow{\scriptstyle f \times f'} \\ \widetilde{\mathrm{Sym}^m X} \times \widetilde{\mathrm{Sym}^m X} & \xrightarrow{j \times id} & \widetilde{\mathrm{Sym}^n X} \times \widetilde{\mathrm{Sym}^m X} \end{array}$$

Then since we have above Cartesian square and f, f' are birational, we have that

$$(j \times id)^* \Gamma' = (j \times id)^* (f \times f')_* (\Gamma)$$

$$= (f' \times f')_* (i \times id)^* (\Gamma) = (f' \times f')_* (\Delta + Y)$$

Here i is the closed embedding of $\mathrm{Sym}^m X$ into $\mathrm{Sym}^n X$, Δ is the diagonal of $\mathrm{Sym}^m X \times \mathrm{Sym}^m X$, and Y is supported on $\mathrm{Sym}^m X \times \mathrm{Sym}^{m-1} X$. Now

$$(f' \times f')_* (\Delta) = \Delta$$

the diagonal of $\widetilde{\mathrm{Sym}^m X} \times \widetilde{\mathrm{Sym}^m X}$ and $(f' \times f')_*(Y)$ is supported on $\widetilde{\mathrm{Sym}^m X} \times \widetilde{\mathrm{Sym}^{m-1} Y}$. The multiplicity of the diagonal of $\widetilde{\mathrm{Sym}^m X} \times \widetilde{\mathrm{Sym}^m X}$ is 1, because the morphism f from $\mathrm{Sym}^n X$ to $\widetilde{\mathrm{Sym}^n X}$ is birational and the multiplicity of Δ of $\mathrm{Sym}^m X \times \mathrm{Sym}^m X$ is 1.

Let ρ be the open embedding of the complement of $\widetilde{\mathrm{Sym}^{m-1} X}$ into $\widetilde{\mathrm{Sym}^m X}$. Then the above computation shows that

$$\rho^* \Gamma'_* j_*(z) = \rho^*(z + z_1) = \rho^*(z)$$

where z_1 is supported on $\widetilde{\mathrm{Sym}^{m-1} X}$. Now consider the following commutative square with the rows exact.

$$\begin{array}{ccccc} CH^*(\widetilde{\mathrm{Sym}^{m-1} X}) & \xrightarrow{j'_*} & CH^*(\widetilde{\mathrm{Sym}^m X}) & \xrightarrow{\rho^*} & CH^*(U) \\ \downarrow & & \downarrow{\scriptstyle j_*} & & \downarrow \\ CH^*(\widetilde{\mathrm{Sym}^{m-1} X}) & \xrightarrow{j''_*} & CH^*(\widetilde{\mathrm{Sym}^n X}) & \longrightarrow & CH^*(V) \end{array}$$

Suppose that $j_*(z) = 0$, then we have $\rho^* \Gamma'_* j_*(z) = 0$, that gives us $\rho^*(z) = 0$. By the localisation exact sequence we have that $z = j'_*(z')$. By the commutativity of the above diagram we have that $j''_*(z') = 0$ and by induction hypothesis we have that $z' = 0$ and hence $z = 0$. So the homomorphism j_* is injective.

□

4. Fano variety of lines on a cubic fourfold and algebraic cycles

In this section we consider the rational cubic fourfolds and the Fano variety of lines on them. It is known [give reference] that the Fano variety is birational to the Hilbert scheme of two points on a K3 surface, provided that the cubic itself is rational. We want to apply the results that we obtained in the previous sections to understand algebraic cycles on the Fano variety of lines of a rational cubic fourfold. More generally when the Fano variety of lines admit of a rational map from the Hilbert scheme of two points on a surface S. The precise theorem is as follows.

Theorem 4.1. *Let X be a cubic fourfold. Let $F(X)$ denote the Fano variety of lines on X. Suppose that there exists a birational map from* $\mathrm{Hilb}^2(S)$,

the Hilbert scheme of two points on a surface S, to $F(X)$. Let $\widetilde{S}$ be the strict transform of S, under the Blow up from $\mathrm{Hilb}^2(S)$ *to* $\mathrm{Sym}^2(S)$*,which intersect the indeterminacy locus of the above bi-rational map transversally. Then we have a codimension* 2 *or codimension* 1 *subvariety D in $F(X)$, (birational to $\widetilde{S}$ or birational to a $\mathbb{P}^1$-bundle over $\widetilde{S}$) such that the kernel of the pushforward $CH_0(D)$ to $CH_0(F(X))$ is torsion.*

Proof. Let us consider the rational map from $\mathrm{Hilb}^2(S)$ to $F(X)$. Let us resolve the indeterminacy of the rational map and we get a regular map

$$\widehat{\mathrm{Hilb}^2(S)} \to F(X)$$

where $\widehat{\mathrm{Hilb}^2(S)}$ denote the Blow up of $\mathrm{Hilb}^2(S)$ along the indeterminacy locus. Suppose that this locus is smooth and intersect a copy of $\widetilde{S}$ transvesally. Then first we prove that the closed embedding $\widehat{S} \to \widehat{\mathrm{Hilb}^2(S)}$ gives an injective push-forward homomorphism at the level of Chow groups of zero cycles, here $\widehat{S}$ is the strict transform of $\widetilde{S}$ under this blow-up. Let f_1 be the natural morphism from $\widehat{\mathrm{Hilb}^2(S)}$ to $\mathrm{Hilb}^2(S)$ and f_1' be the restriction of it to $\widehat{S}$. Consider the correspondence Γ'' on $\widehat{\mathrm{Hilb}^2(S)} \times \widehat{S}$ given by $(f_1 \times f_1')^*(\Gamma')$. Let $\hat{j}$ denote the closed embedding of $\widehat{S}$ into $\widehat{\mathrm{Hilb}^2(S)}$. Then as previous we have that $\Gamma''_* \hat{j}_*$ is induced by $(\hat{j} \times id)^*(\Gamma'')$, which is equal to

$$(\hat{j} \times id)^*(f_1 \times f_1')^* \Gamma'$$

which is

$$(f_1' \times f_1')^*(\Delta + Y_1 + E_1)$$

where E_1 is the intersection of the exceptional locus of the blow up $\mathrm{Hilb}^2(S) \to \mathrm{Sym}^2 S$ with $\widetilde{S}$ and Y_1 is supported on $\widetilde{S} \times \mathbb{P}^1$. Then we have

$$(f_1' \times f_1')^*(\Delta + Y_1 + E_1) = \Delta + \hat{E}_2 + \hat{E}_1 + \hat{Y}_1 \,.$$

Here $\hat{E}_2$ is the intersection of the exceptional locus $\hat{E}$ of the blow up $\widehat{\mathrm{Hilb}^2(S)} \to \mathrm{Hilb}^2(S)$ with $\hat{S}$. Then consider the open embedding of the complement of $\hat{Y}$ in $\hat{S}$ (where $\hat{Y}$ is a $\mathbb{P}^1$-bundle over $\mathbb{P}^1$, or a rational curve). Call it ρ. We have by Chow moving lemma,

$$\rho^* \Gamma''_* \hat{j}_*(z) = \rho^*(z + z_1) = \rho^*(z) \,.$$

Here z_1 is supported on A. Consider the following commutative diagram where the rows are exact.

$$\begin{array}{ccccc} CH^*(\hat{Y}) & \xrightarrow{j'_*} & CH^*(\hat{S}) & \xrightarrow{\rho^*} & CH^*(U) \\ \downarrow & & \downarrow{\scriptstyle \hat{j}_*} & & \downarrow \\ CH^*(\hat{Y}) & \xrightarrow{j''_*} & CH^*(\widehat{\mathrm{Hilb}^2(S)}) & \longrightarrow & CH^*(V) \end{array}$$

Suppose that $\hat{j}_*(z) = 0$, then we have $\rho^*\Gamma''_*\hat{j}_*(z) = 0$, which gives $\rho^*(z) = 0$. So there exists z' such that $j_*(z') = z$. The map from $CH_0(\hat{Y}) \to CH_0(\widehat{\mathrm{Hilb}^2 S})$ is injective. So $z' = 0$, hence $z = 0$. Now we have a regular birational (generically finite) map from $\widehat{\mathrm{Hilb}^2(S)}$ to $F(X)$ and let D denote the codimension 2 image of $\hat{S}$. Then we prove that the kernel of the push-forward homomorphism from $CH_0(D)$ to $CH_0(F(X))$ is torsion. So let h be the closed embedding of D into $F(X)$ and g the map from $\widehat{\mathrm{Hilb}^2(S)}$ to $F(X)$ and g' its restriction to $\hat{S}$. Then consider the correspondence Γ''' to be $(g \times g')_*(\Gamma'')$. The homomorphism $\Gamma'''_* h_*$ is induced by the cycle $(h \times id)^*(\Gamma''')$, which is

$$(h \times id)^*(g \times g')_*(\Gamma'')$$

and is equal to the fundamental cycle [Fu][1.5] associated to (since g is birational and generically finite)

$$(g' \times g')_(\hat{j} \times id)^{-1}(\Gamma'')$$

that is equal to

$$(d\Delta + (g' \times g')_*(\hat{E}_2 + \hat{E}_1 + \hat{Y}))$$

where the later is supported on $D \times V$ where V is the image of the $\mathbb{P}^1$ bundle $\hat{Y}$ or on $D \times C$ where C is a rational curve (possibly singular) on D. So let ρ be the open embedding of the complement of $g'(\hat{Y})$ in D. Then as previous we have that

$$\rho^*\Gamma'''_* h_*(z) = \rho^*(dz + z_1)$$

where z_1 is supported on $g'(\hat{Y})$. Then consider the following commutative diagram with the rows exact.

$$\begin{array}{ccccc} CH^*(g'(\hat{Y})) & \xrightarrow{h'_*} & CH^*(D) & \xrightarrow{\rho^*} & CH^*(U) \\ \downarrow & & \downarrow{\scriptstyle h_*} & & \downarrow \\ CH^*(g'(\hat{Y})) & \xrightarrow{j''_*} & CH^*(F(X)) & \longrightarrow & CH^*(V) \end{array}$$

Then if $h_*(z) = 0$, it would follow that $d(\rho^*(z)) = 0$. So we have $h'_*(z') = dz$. Now $CH_0(g'(\hat{Y})) \to CH_0(F(X))$ is injective as V is rational or $CH_0(g'\hat{Y})$ is torsion since C is rational, tell us that $z' = 0$ or $d'z' = 0$ so $dz = 0$ or $dd'z = 0$. So the kernel of h_* is torsion. □

Now all this techniques can be applicable in the following setup. Let us embed $\mathrm{Sym}^2 S$ into some projective space. Let C be a smooth hyperplane section of S inside $\mathrm{Sym}^2 S$ such that C intersects the diagonal transversally (or do not intersect). Then consider the strict transform of $\widetilde{C}$ inside $\widetilde{\mathrm{Sym}^2 S} = \mathrm{Hilb}^2(S)$, that gives us push-forward homomorphism at the level of Chow group of zero cycles, which has torsion kernel. When we blow up $\mathrm{Hilb}^2 S$, then we get the strict transform $\hat{C}$, which is again inducing an push-forward at the level of Chow groups from $CH_0(\hat{C})$ to $CH_0(\widehat{\mathrm{Hilb}^2(S)})$ having torsion kernel. Then we push-down everything to $F(X)$, we get that the image of $\hat{C}$, inside $F(X)$, say C', is such that the push-forward $CH_0(C') \to CH_0(F(X))$ has torsion kernel.

Let the ground field k be such that the affine line is not a zero divisor in the Grothendieck ring of varieties over k. In this case the assumption of the above theorem is true, that is there is a birational map from $\mathrm{Hilb}^2(X)$ to $F(X)$, where X is a rational cubic fourfold and we arrive at the following result.

Theorem 4.2. *If the cubic is rational and if the indeterminacy locus of the rational map* $\mathrm{Hilb}^2 S \to F(X)$ *is smooth, then there exists a projective curve, or a surface (admitting a finite map from a ruled surface) in* $F(X)$ *such that the kernel of the push-forward (induced by the closed embedding of the curve or the surface inside* $F(X)$*) at the level of Chow groups of zero cycles in torsion. Contra-positively we can say that if for all projective curves and projective surfaces in* $F(X)$*, the kernel of the push-forward induced by*

the closed embedding of the curve or the surface has non-torsion elements, then the cubic is non-rational.

REFERENCES

[B] K.Banerjee,*Blow ups and base changes of symmetric powers and Chow groups*, https://sites.google.com/site/kalyanmath484/publications, 2015.

[BI] K.Banerjee, Jaya NN Iyer, *On the kernel of the push-forward homomorphism between Chow groups*, arxiv::1504.07887v1,2015.

[BG] K. Banerjee and V. Guletskii, *Rational equivalence for line configurations on cubic hypersurfaces in* $\mathbb{P}^5$, arXiv:1405.6430v1, 2014.

[Collino] A. Collino, *The rational equivalence ring of symmetric product of curves*, Illinois journal of mathematics, 19, no.4, 567-583, 1975.

[Fu] W. Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 2. Springer-Verlag, Berlin, 1984.

[GS] S.Galkin, E.Shinder, *The Fano variety of lines and the rationality problem of a cubic hypersurface*, https://arxiv.org/abs/1405.5154

[Ha] B.Hassett, *Special cubic fourfolds*, Compositio Math. 120 No.1, 1-23. 2000.

INDIAN STATISTICAL INSTITUTE, BANGALORE CENTER, BANGALORE 560059
E-mail address: kalyanb_vs@isibang.ac.in